

The community structure of macroscopic basidiomycetes (Fungi) in Brazilian mangroves influenced by temporal and spatial variations

Georgina Santos Nogueira-Melo¹, Paulo Jorge Parreira Santos² & Tatiana Baptista Gibertoni¹

1. Departamento de Micologia, Universidade Federal de Pernambuco, Av. Nelson Chaves s/n, CEP 50760-420, Recife, PE, Brazil; geomycota@gmail.com, tatiana.gibertoni@pq.cnpq.br
2. Departamento de Zoologia, Universidade Federal de Pernambuco, Av. Nelson Chaves s/n, CEP 50760-420, Recife, PE, Brazil; pjp.santos@gmail.com

Received 12-XII-2013. Corrected 22-V-2014. Accepted 24-VI-2014.

Abstract: Mangroves are transitional ecosystems between terrestrial and marine environments, and are distinguished by a high abundance of animals, plants, and fungi. Although macrofungi occur in different types of habitat, including mangroves, little is known about their community structure and dynamic. Therefore the aim of this study was to analyze the diversity of macrofungi in a number of Brazilian mangroves, and the relationship between such diversity, precipitation and area of collection. A total of 32 field trips were undertaken from 2009 to 2010, and macrofungi were studied in four 250×40m transects: Timbó and Santa Cruz Channel on the Northern coast, and Maracaípe and Ariquindá on the Southern coast. All basidiomata found along the transects were placed in paper bags, air-dried and identified using existing literature. It was found that Northern areas predominantly featured *Avicennia schaueriana* mangroves, while *Rhizophora mangle* dominated in Southern transects. A total of 275 specimens were collected, and 33 species, 28 genera, 14 families and six orders were represented. Overall abundance and species richness did not vary significantly among areas, but varied according to time, being higher during the rainy season. Subtle differences in composition were observed over time and between areas, probably due to variations in plant species occurrence. Further studies with collections during months of greater precipitation in transects dominated by different mangrove species of the same ecosystem are suggested to assess the overall diversity of mycobiota in these ecosystems. Rev. Biol. Trop. 62 (4): 1587-1595. Epub 2014 December 01.

Key words: Agaricomycetes, diversity, ecological interactions, estuaries, fungi.

Mangroves are transitional ecosystems between terrestrial and marine environments and are important for the maintenance of biodiversity and water quality, sediment fixing, and the supply of primary production to the surroundings (Cintrón & Schaeffer-Novelli, 1980). Mangroves are known for their particular vegetation and, while not being species-rich, are distinguished by a high abundance of animals, plants, and fungi (Silva, Bernini & Carmo, 2005).

Macrofungi occur in different habitat types, including mangroves, and are found with higher frequency and diversity in tropical forests. Due to their saprotrophic, parasitic,

or symbiotic lifestyles, they play an essential role in the ecological balance of forest areas. Macroscopic basidiomycetes (a type of fungi known as mushrooms, bracket fungi, earth-stars, among others) can be found on living or dead wood, in soil, in mycorrhizal or lichen associations, or as plant parasites (Alexopoulos, Mims & Blackwell, 1996; Kendrick, 2000; Deacon, 2006; Webster & Weber, 2007).

Despite the importance of decaying wood fungi, little is known about their community structure and dynamic in mangroves. Information of the distribution patterns and the factors that control or limit their diversity in such environments in tropical areas is scarce (Lindblad,

2000; Lindblad, 2001; Gilbert & Sousa, 2002; Urcelay & Robledo, 2004; Robledo, Urcelay, Domínguez & Rajchenberg, 2006; Gibertoni, Santos & Cavalcanti, 2007; Gibertoni, 2008; Gilbert, Gorospe & Ryvardeen, 2008; Drechsler-Santos, Santos, Gibertoni & Cavalcanti, 2010; Robledo & Renison, 2010; Trierweiler-Pereira, Santos & Baseia, 2013). Of these, only two studies were undertaken in mangroves. Gilbert & Sousa (2002) observed that macrofungi occur more frequently in a particular type of substrate, while Gilbert et al. (2008) observed the host and habitat preferences of polypores in three well-defined, floristically distinct, tropical wetlands, which included mangroves. In the latter case, the habitat preference was related to specificity for a particular host. In Brazil only species checklists and descriptions are available (Sotão, Bononi, & Figueiredo, 1991; Almeida-Filho, Bueno & Bononi, 1993; Gugliotta & Capelari, 1995; Campos & Cavalcanti, 2000; Campos, Sotão, Cavalcanti & Luz, 2005; Baltazar, Trierweiler-Pereira & Loguercio-Leite, 2009; Nogueira-Melo, Ryvardeen & Gibertoni, 2011; Nogueira-Melo et al., 2012)

The aim of the present study was to determine the structure and describe the specific composition of a wood-decaying fungi community through the occurrence of basidiomata in four different mangroves areas in Brazil. Spatial variations between areas and temporal variations between the two main seasons of the year were also described. The data was used to test the hypothesis that diversity of fungi is directly affected by both rainfall and area of collection.

MATERIALS AND METHODS

The study was conducted in four mangrove areas, belonging to four estuaries in the state of Pernambuco in the Northeast of Brazil. Two were along the Northern coast: mangrove from the Timbó river (T) (07°51'24"8" S - 34°50'32"7" W) and mangrove from the Santa Cruz Chanel (S) (07°46'52"6" S - 34°52'53"3" W), where *Avicennia schaueriana* predominates; and two along the Southern

coast: mangrove from the Maracaípe river (M) (08°32'22"8" S - 35°00'29"1" W) and mangrove from the Ariquindá river (A) (08°41'20"8" S - 35°06'6"6" W), where *Rhizophora mangle* predominates. Northern coast mangroves were surrounded by urban areas, while those from the Southern coast were distant from such areas (Montes, Macêdo & Koenig, 2002; Mendonça & Almeida-Cortez, 2007). The Northern mangroves are approximately 45km apart, while those from the Southern coast are 40km apart. The Northern and Southern mangroves are around 100km apart.

Three typical mangrove tree species were frequently found in the study areas: *Avicennia schaueriana* Stapf. and Leech (black mangrove), *Laguncularia racemosa* (L.) Gaertn (white mangrove), and *Rhizophora mangle* L. (red mangrove). In the studied forests, these species are differentially distributed according to distance from the water's edge, but their distributions overlap to varying degrees (Cintrón & Schaeffer-Novelli, 1980; Schaeffer-Novelli, Cintrón-Molero, Andaime & Camargo, 1990).

The climate in these areas is defined as tropical humid (Am by Köepen classification), with high monthly and low annual temperature amplitude, an average annual temperature of 21°C and annual precipitation greater than 1 500mm (Peel, Finlayson & McMahon, 2007).

In order to study fungi in these areas, one transect of 250×40m (10 000m²; 20m to the right and 20m to the left of the transect), starting in the estuary and roughly following a line parallel to the water channel, was established in each mangrove, using Global Position System (GPS). A collection trip was undertaken monthly in each of the four areas for eight months (Apr, May, Jun, Jul, Aug, Sep, 2009; Jan and Mar 2010), totaling 32 field trips (16 along each coast).

All basidiomata found along transects were placed in paper bags and collection data (position, date and substrate) was recorded. The material was air-dried and identified with help of existing literature (Ryvardeen & Johansen, 1980; Gilbertson & Ryvardeen, 1986; 1987; Hjortstam, Larsson & Ryvardeen, 1987; 1988;

Ryvarden, 1991; Boidin, Lanquetin & Gilles, 1997; Boidin & Gilles, 2000; Núñez & Ryvarden, 2000; 2001; Ryvarden, 2004; Bernicchia, 2005). Nomenclature from the CABI online database (www.indexfungorum.org) was used. Identified material in good condition was incorporated into the URM Herbarium, of the Department of Mycology of the Universidade Federal de Pernambuco, and duplicates were sent to the O Herbarium of the University of Oslo. Samples of the same species collected in the same area in the same substrate and on the same date were identified as one combined voucher specimen.

Species abundance values were represented by the number of occurrences of specimens/individuals in each substrate per transect (ind./10 000m²). One specimen/individual may be represented by several basidiomata.

Statistical analysis was performed in accordance with the methods proposed by Clarke & Warwick (1994) using the PRIMER[®] 6 (Plymouth Routines in Multivariate Ecological Research) software.

Two-way variance analysis (ANOVA) was used to determine significant differences for abundance and richness between both mangroves and precipitation periods (dry and wet), and two-way similarity analysis (ANOSIM) was used to determine significant differences in the macrofungi community structure between the four mangroves and sampling seasons. Bartlett's test was used to determine the homogeneity of variances and values were transformed into Neperian logarithms if necessary.

The Bray–Curtis index was applied to standardized data to assess the similarity between replicates. Multi-dimensional scaling (MDS) was used to represent the similarity matrix.

The BioEstat 5.0 method of binomial probability distribution was used to evaluate spatial distribution and verify if there was predominance of some fungal species in any of the mangroves. Species considered not to be rare were used in the analysis and the areas were tested separately. In the present study, rare species were those whose frequency of occurrence was $0.5 < F \leq 1.5\%$, using the formula:

$F = n \times 100 / N$, where n = number of specimens from one species and N = total number of specimens (Lindblad, 2000; Schnittler & Stephenson, 2000). The probability value used (P) was that of species with higher incidence (Ayres, Ayres, Ayres & Santos, 2007).

The normal approximation to the binomial test was used to verify the differences among plant proportions in areas using BioEstat 5.0 software. Since six paired comparisons were made, the significance level was set at $p < 0.01$ for this analysis. If not stated, the level of significance was set at $p < 0.05$ for all other analyses.

Abiotic factors considered in this study were temperature and rainfall, data of which was provided by the Instituto de Tecnologia de Pernambuco - ITEP/Agência Pernambucana de Água e Clima (APAC). Dry months (Sep 2009, Jan and Mar 2010) were considered those in which the total value of precipitation was below the lower limit of the confidence interval (95%) for the average annual monthly precipitation over the last 10 years.

RESULTS

A total of 275 specimens were analyzed, representing 33 species, 28 genera, 14 families and six orders. Of the 14 families, four presented more than 15 records and together constituted 70% of specimens collected: Gloeophyllaceae, with 55 records, was represented only by *Gloeophyllum striatum*; Hymenochaetaeaceae, which had 25 occurrences of two genera and four species; Polyporaceae, which had 95 occurrences corresponding to nine species and eight genera; and Schizoporaceae, which had 26 records and was represented exclusively by *Schizopora paradoxa* (Table 1).

From the 275 collected specimens, 81 occurred in S, 66 in T, 73 in M and 55 in A. The Santa Cruz Chanel presented the highest number of genera and species (18 and 20, respectively), followed by T (15 genera, 16 species), M (12, 16) and A (12, 13). Although richness and abundance varied, they did not differ significantly between areas [$F_{(3,21)} = 3.324$,

TABLE I

Wood decaying fungi collected in four Brazilian mangroves (A=Ariquindá; M=Maracaípe; S=Santa Cruz Chanel; T=Timbó) and occurrence by substrate (As=*Avicennia schaueriana*; Lr=*Laguncularia racemosa*; Rm=*Rhizophora mangle*)

Species	A		M		S		T	
	Rm	Lr	Rm	As	Rm	As	Rm	As
CORTICIACEAE Herter								
<i>Punctularia strigosozonata</i> (Scheinw.) P.H.B. Talbot						x		
DACRYMYCETACEAE J. Schröt.								
<i>Cerinomyces aculeatus</i> N. Maekawa					x			
FOMITOPSISIDACEAE Jülich								
<i>Fomitopsis nivos</i> (Berk.) Gilb. & Ryvarden	x					x	x	
GLOEOPHYLLACEAE Jülich								
<i>Gloeophyllum striatum</i> (Sw.) Murrill	x		x		x		x	
HYMENOCHEAETACEAE Donk								
<i>Fuscoporia contigua</i> (Pers.) G. Cunn.						x		x
<i>Phellinus</i> cf. <i>rhytiphloeus</i> (Mont.) Ryvarden							x	x
<i>Phellinus gilvus</i> (Schwein.) Pat.	x		x	x	x	x	x	x
<i>Phellinus mangrovicus</i> (Imazeki) Imazeki			x					
<i>Phellinus rimosus</i> (Berk.) Pilát	x		x					
LACHNOCLADIACEAE D.A. Reid								
<i>Asterostroma cervicolor</i> (Berk. & M.A. Curtis) Massee					x		x	
MERULIACEAE Rea								
<i>Cerocorticium molle</i> (Berk. & M.A. Curtis) Jülich			x				x	x
<i>Gloeoporus dichrous</i> (Fr.) Bres.					x			
<i>Hyphoderma iguazuense</i> Hjortstam & Ryvarden	x	x					x	x
<i>Hyphoderma</i> sp.					x			
PENIOPHORACEAE Lotsy								
<i>Gloeocystidiopsis</i> cf. <i>salmonia</i> (Burt) Boidin, Lanq. & Gilles			x					x
PHANEROCHAETACEAE Jülich								
<i>Ceriporia spissa</i> (Schwein. ex Fr.) Rajchenb.			x					
<i>Phanerochaete australis</i> Jülich					x			
<i>Phlebiopsis ravenelii</i> (Cooke) Hjortstam			x	x				
POLYPORACEAE Fr. ex Corda								
<i>Coriolopsis hostmanii</i> (Berk.) Ryvarden						x	x	x
<i>Hexagonia hydnooides</i> (Sw.) M. Fidalgo						x	x	x
<i>Hjortstamia amethystea</i> (Hjortstam & Ryvarden) Boidin & Gilles	x		x			x		x
<i>Lentinus bertieri</i> (Fr.) Fr.		x						
<i>Lopharia</i> sp.					x			
<i>Loweporus tephroporus</i> (Mont.) Ryvarden						x		
<i>Perenniporia detrita</i> (Berk.) Ryvarden	x		x		x	x		
<i>Perenniporia guyanensis</i> Decock & Ryvarden			x	x				
<i>Pycnoporus sanguineus</i> (L.) Murrill						x		
<i>Trichaptum bifforme</i> (Fr.) Ryvarden		x	x		x	x	x	x
SCHIZOPHYLLACEAE Quéf.								
<i>Schizophyllum commune</i> Fr.			x		x		x	x
SCHIZOPORACEAE Jülich								
<i>Schizopora paradoxa</i> (Schrad.) Donk	x		x		x	x	x	x
STEREACEAE Pilát								
<i>Gloeocystidiellum triste</i> Hjortstam & Ryvarden			x			x	x	
<i>Gleodontia discolor</i> (Berk. & M.A. Curtis) Boidin		x	x			x		
TRICHOLOMATACEAE R. Heim ex Pouzar								
<i>Resupinatus poriaeformis</i> (Pers.) Thorn, Moncalvo & Redhead								x




Fig. 1. Multidimensional scaling graphic of studied areas: M=Maracaípe; A=Ariquindá; T=Timbó, and S=Santa Cruz Chanel. Resemblance: S17 Bray Curtis similarity. 2D Stress: 0.19.

$p=0.015$; $F_{(3;21)}=1.060$, $p=0.388$, respectively]. ANOSIM revealed subtle differences in the proportion of species of the fungal community among the four mangrove areas ($R_{\text{global}}=0.384$, number of permutations=999, $p=0.001$), and between the northern (S and T) and southern mangroves (A and M) ($R_{\text{global}}=0.33$, number of permutations=999, $p=0.001$) (Fig. 1).

Of the 33 species, four were common to all four areas: *G. striatum*, *P. gilvus*, *S. paradoxa* and *T. biforme*. Significantly different rates of occurrence according to area were found for nine of the 11 fungal species (82%) which were sufficiently abundant for statistical testing (Table 2, in bold). *Corioloopsis hostmanii*, *P. gilvus* and *T. biforme* were predominant in S; *H. hydnoides* and *H. iguazuense* in T; *P. guyanensis* and *G. striatum* in M, and *S. paradoxa* in A, while the values of *H. amethystea* and *P. detrita* were not different from random occurrence levels (Table 2).

The studied areas differed in proportions of mangrove species. Northern coastal

transects showed predominance of *A. schaueriana*, while Southern coastal transects presented mostly *R. mangle* ($M \times A: Z=1.043$, $p=0.297$; $M \times S: Z=7.677$, $p<0.0001$; $M \times T: Z=9.065$, $p<0.0001$; $A \times S: Z=6.680$, $p<0.0001$; $A \times T: Z=8.072$, $p<0.0001$; $S \times T: Z=2.220$, $p=0.0264$).

Average rainfall in the four areas from April 2009 to March 2010 was 174.13mm, varying monthly from 0mm to 543.2mm. The mean temperature was 25.17°C. The driest month was October (11.6mm) and the wettest month was April (409.85mm). January, February, March, September, October, November and December were considered to be the dry season, while April, May, June, July and August represented the rainy season (Fig. 2).

Of the 275 collected specimens, 30 (=15 species) were found in April, 41 (=17) in May, 53 (=20) in June, 40 (=14) in July, 31 (=16) in August, 35 (=16) in September, 32 (=11) in December and 13 (=six) in March. There was a significant difference in sample richness

TABLE 2

Wood decaying fungi collected in four Brazilian mangroves of sufficient abundance for analysis of predominance by area (S=Santa Cruz Chanel; M=Maracaípe; T=Timbó and A=Ariquindá)

Species	S	T	M	A	p
<i>Trichaptum biforme</i> (Fr.) Ryvarden	25	7	1	8	0.0001
<i>Corioliopsis hostmannii</i> (Berk.) Ryvarden	10	1	0	0	0.001
<i>Phellinus gilvus</i> (Schwein.) Pat.	9	5	1	2	0.0028
<i>Hexagonia hydnoides</i> (Sw.) M. Fidalgo	4	9	0	0	0.0018
<i>Cerocorticium molle</i> (Berk. & M.A. Curtis) Jülich	0	8	1	3	0.0039
<i>Hyphoderma iguazuense</i> Hjortstam & Ryvarden	0	8	0	2	0.0014
<i>Gloeophyllum striatum</i> (Sw.) Murrill	5	11	32	17	0.0029
<i>Perenniporia guyanensis</i> Decock & Ryvarden	0	0	9	0	0.0018
<i>Schizophora paradoxa</i> (Schrad.) Donk	5	5	4	12	0.0076
<i>Perenniporia detrita</i> (Berk.) Ryvarden	1	0	6	3	0.0592
<i>Lopharia amethystea</i> (Hjortstam & Ryvarden) A.L.Weden	3	2	2	1	0.1478
Number of trees	77	82	117	92	




Fig. 2. Temperature and precipitation in the studied areas from April 2009 to March 2010. M=Maracaípe; A=Ariquindá; T=Timbó, and S=Santa Cruz Chanel.

according to time between June and March [$F_{(3;21)}=3.324$, $p=0.015$]. Additionally, there were temporal differences in abundance among the collection periods: May, June, July and September yielded more specimens than March [$(F_{(3;21)}=4.055$, $p=0.006)$]. A slight difference in species composition was also observed ($R_{\text{global}}=0.141$, number of permutations=999, $p=0.044$) when occurrence of fungi in the dry and rainy season were compared.

DISCUSSION

The present study showed that the relative proportion of macroscopic basidiomycete

species was different among the four studied areas, and between the Northern and Southern mangroves. Although the ANOSIM R_{global} value found a statistical significant distinction among areas ($R_{\text{global}}=0.384$ and 0.33 , respectively), according to Clarke & Warwick (1994), genuinely different communities would result in R_{global} values above 0.5.

Therefore, the collected species belong to the same community (macrofungi inhabiting mangroves). However, the differential occurrence of about one-third of the 33 collected fungal species in the areas may have contributed to the observed distinction, although subtle, between areas.

Furthermore, in transects of the Northern mangroves, *A. schaueriana* was predominant, while *R. mangle* was predominant in transects of the Southern mangroves. In estuaries, salinity and/or proximity to the sea influenced the distribution of mangrove plant species (Cintrón & Schaeffer-Novelli, 1980; Schaeffer-Novelli et al., 1990). Additionally, the Northern mangroves were surrounded by urban areas, while the Southern mangroves were distant from such areas (Montes et al., 2002; Mendonça & Almeida-Cortez, 2007), and thus have less anthropic impacts. Hyde (1989) described the negative effects of hydrocarbon spillage on microfungi diversity in mangroves in Brunei, while Yamashita et al. (2008) described the negative effects of agriculture and isolated patches of natural forest on aphylloraceous diversity in Malaysia. However, the hypothesis that anthropization influences fungal communities was not tested in the present work. Hence, the distribution of the fungal species may reflect the vegetation in the studied transects.

In the present study, the abundance and richness of wood-decaying basidiomycetes (and to a lesser degree composition) were affected by precipitation: more specimens were collected in the rainy season and fewer during the dry season. The influence of precipitation on the occurrence of these macrofungi has also been reported in other ecosystems in Brazil. Gibertoni et al. (2007) observed that the former aphylloroid fungi were more frequently collected in the dry season, after peaks of rainfall in the Atlantic Forest. In Amazonia, Gibertoni (2008) found that the composition of polyporoid fungi species was different in the rainy than in the dry season. This indicates that rainfall is a factor that influences the occurrence of species and specimens.

However, Drechsler-Santos et al. (2010) observed in caatinga (semi-arid region), that rainfall did not influence the occurrence of species of Hymenochaetaceae, mostly found in living hosts. This is probably because they do not depend on environmental humidity as they are adapted to moisture hosts, according to Boddy & Rayner (1983).

Recently, Trierweiler-Pereira et al. (2013) reported no significant differences in composition and species richness of epigeous Gasteromycetes according to collecting seasons in the Atlantic Forest, probably due to ephemeral basidiomata, which may not be observed, even during the rainy season.

The results of the present study support the tested hypothesis that diversity of macrofungi occurring in mangroves is directly affected by both rainfall and area of collection. We recommend further studies should collect samples during the higher precipitation months and from transects dominated by different mangrove plants in the same area in order to identify the overall diversity of the mycobiota in this ecosystem.

ACKNOWLEDGMENTS

The authors would like to thank their laboratory colleagues for support during the field trips; Leif Ryvarden and Karl-Henrik Larsson for species identification, Robert Lücking and Michael Finnie for English improvement and valuable suggestions, Roger Melo and Lucas Cavalcanti for help with figures. This research was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, master scholarship of GSNM) and by FACEPE (APQ 0444-2.03/08).

RESUMEN

La estructura de la comunidad de basidiomicetos macroscópicos (Fungi) en los manglares brasileños influenciada por las variaciones temporales y espaciales. Los manglares son ecosistemas de transición entre los ambientes terrestres y marinos, y se distinguen por la gran abundancia de animales, plantas y hongos. Aunque los macrohongos se encuentran en diferentes tipos de hábitat, incluidos los manglares, poco se sabe acerca de la estructura de su comunidad y dinámica. Por lo tanto, el objetivo de este estudio fue analizar la diversidad de macrohongos en los manglares de Brasil y su relación con la precipitación y área de recolección. Se realizaron un total de 32 salidas de campo entre 2009 y 2010, y los macrohongos fueron estudiados en cuatro transectos de 250×40m: Timbó y Canal de Santa Cruz en la costa norte y Maracaípe y Ariquindá en la costa sur. Todos los basidiomas encontrados

a lo largo de los transectos se colocaron deshidratados en bolsas de papel, y se identificaron con ayuda de la literatura preexistente. Se encontró que las zonas del norte predominantemente presentaron *Avicennia schaueriana*, mientras *Rhizophora mangle* domina en transectos del sur. Se recolectaron un total de 275 especímenes y 33 especies, 28 géneros, 14 familias y seis órdenes estuvieron representados. Abundancia y riqueza de especies en general no varió significativamente entre las áreas, pero sí varió en el tiempo, siendo mayor durante la estación lluviosa. Se observaron diferencias sutiles en la composición a través del tiempo y entre áreas, probablemente debido a las variaciones en la presencia de las especies de plantas. Otros estudios con recolectas durante los meses de mayor precipitación en transectos dominados por diferentes plantas de manglar en el mismo ecosistema son deseables para acceder a la diversidad de la microbiota.

Palabras clave: Agaricomycetes, diversidad, interacciones ecológicas, estuarios, hongos.

REFERENCES

- Alexopoulos, C. J., Mims, C. W., & Blackwell, M. (1996). *Introductory Mycology* (4th ed.). New York: John Wiley and Sons Press.
- Almeida-Filho, O. M., Bueno, R., & Bononi, V. L. R. (1993). Algumas espécies de fungos basidiomicetos dos manguezais do Estado de São Paulo. *Hoehnea*, 20(1/2), 87-92.
- Ayres, M., Ayres, J. R. M., Ayres, D. L., & Santos, A. A. S. (2007). *BioEstat: Aplicações estatísticas nas áreas de ciências biomédicas* (5th ed.). Belém: Guanabara.
- Baltazar, J. M., Trierweiler-Pereira, L., & Loguercio-Leite, C. (2009). A checklist of xylophilous basidiomycetes (Basidiomycota) in mangroves. *Mycotaxon*, 107, 221-224.
- Bernicchia, A. (2005). *Polyporaceae s.l. Fungi europaei 10*. Alassio: Candusso.
- Boddy, L. & Rayner, A. D. M. (1983). Origins of decay in living deciduous trees: the role of moisture content and a re-appraisal of the expanded concept of tree decay. *New Phytology*, 94, 623-641.
- Boidin, J., Lanquetin, P., & Gilles, G. (1997). Le genre *Gleocystidiellum sensu lato* (Basidiomycotina). *Bulletin trimestriel de la Société mycologique de France*, 113.
- Boidin, J. & Gilles, G. (2000). Basidiomycètes Aphyllophorales de l'île de la Réunion XXI-suite. *Mycotaxon*, 75, 357-387.
- Campos, E. L. & Cavalcanti, M. A. Q. (2000). Primeira ocorrência de *Phellinus mangrovicus* (Imaz.) Imaz. para o Brasil. *Acta botânica Brasileira*, 14(3), 263-265.
- Campos, E. L., Sotão, H. M. P., Cavalcanti, M. A. Q., & Luz, A. B. (2005). *Basidiomycetes* de manguezais da APA de Algodual-Maiandeuá, Pará, Brasil. *Boletim do Museu Paraense. Emílio Goeldi*, 1(1), 97-102.
- Cintrón, G. & Schaeffer-Novelly, Y. (1980). *Introducción a la ecología del manglar*. São Paulo: Instituto Oceanográfico USP.
- Clarke, K. R. & Warwick, R. M. (1994). *Change in marine communities: an approach to statistical analysis and interpretation*. Plymouth: Plymouth Marine Laboratory.
- Deacon, J. W. (2006). *Fungal biology* (4th ed.). Massachusetts: Blackwell Publishing.
- Drechsler-Santos, E. R., Santos, P. J. P., Gibertoni, T. B., & Cavalcanti, M. A. Q. (2010). Ecological aspects of Hymenochaetaceae in an area of Caatinga (semi-arid) in Northeast Brazil. *Fungal Diversity*, 42, 71-78.
- Gibertoni, T. B., Santos, P. J. P., & Cavalcanti, M. A. Q. (2007). Ecological aspects of *Aphyllphorales* in the Atlantic Rain Forest in Northeast Brazil. *Fungal Diversity*, 25, 49-67.
- Gibertoni, T. B. (2008). Polyporoid fungi (*Agaricomycetes, Basidiomycota*) in the Estação Científica Ferreira Penna (State of Pará, Brazilian Amazonia): diversity and ecological aspects. *Scientifica Acta*, 2, 70-74.
- Gilbert, G. S. & Sousa, W. P. (2002). Host specialization among wood-decay polypore fungi in a Caribbean mangrove forest. *Biotropica*, 34(3), 396-404.
- Gilbert, G. S., Gorospe, J., & Ryvarden, L. (2008). Host and habitat preferences of polypore fungi in Micronesian tropical flooded forests. *Mycological Research*, 112(6), 674-680.
- Gilbertson, R. L. & Ryvarden, L. (1986). *North American Polypores*. (Vol. 1). Oslo: Fungiflora.
- Gilbertson, R. L. & Ryvarden, L. (1987). *North American Polypores*. (Vol. 1). Oslo: Fungiflora.
- Gugliotta, A. M. & Capelari, M. (1995). *Polyporaceae* from Ilha do Cardoso, SP, Brazil. *Mycotaxon*, 56, 107-113.
- Hjortstam, K., Larsson, K. H., & Ryvarden, L. (1987). *The Corticiaceae of North Europe*. (Vol. 1). Oslo: Fungiflora.
- Hjortstam, K., Larsson, K. H., & Ryvarden, L. (1988). *The Corticiaceae of North Europe* (Vol. 2). Oslo: Fungiflora.
- Hyde, K. D. (1989). Ecology of tropical marine fungi. *Hydrobiologia*, 178, 199-208.
- Kendrick, B. (2000). *The fifth kingdom* (2nd ed.). Newburyport: Focus Information Group.
- Lindblad, I. (2000). Host specificity of some wood-inhabiting fungi in a tropical forest. *Mycologia*, 92(3), 399-405.

- Lindblad, I. (2001). Diversity of poroid and some corticoid wood-inhabiting fungi along the rainfall gradient in tropical forests, Costa Rica. *Journal of Tropical Ecology*, 17, 353-369.
- Mendonça, I. V. S. & Almeida-Cortez, J. S. (2007). Caracterização da galha induzida por ácaro em *Laguncularia racemosa* (L.) Gaertn (Combretaceae). *Biota Neotropica*, 7(3), 163-170.
- Montes, M. J. F., Macêdo S. J., & Koenig, M. L. (2002). N: Si: P Atomic Ratio in the Santa Cruz Channel, Itamaracá- PE (Northeast Brazil): a Nyctemeral Variation. Brazilian. *Archives of Biology and Technology*, 45(2), 115-124.
- Nogueira-Melo, G. S., Ryvarden, L., & Gibertoni, T. B. (2011). First record of *Resupinatus poriaeformis* (Agaricomycetes) from South America. *Mycotaxon*, 117, 423-427.
- Nogueira-Melo, G. S., Medeiros, P. S., Gomes-Silva, A. C., Ryvarden, L., Sothão, H. M. P., & Gibertoni, T. B. (2012). *Corioloopsis psila* comb. nov. (Agaricomycetes) and two new *Corioloopsis* records for Brazil. *Mycotaxon*, 120, 223-230.
- Núñez, M. & Ryvarden, L. (2000). East Asian Polypores Vol. 1. *Ganodermataceae* and *Hymenochaetaceae*. *Synopsis Fungorum*, 13, 1-168.
- Núñez, M. & Ryvarden, L. (2001). East Asian Polypores. Vol. 2. *Polporaceae sensu lato*. *Synopsis Fungorum*, 14, 169-522.
- Peel, M. C., Finlayson, B. L., & McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences*, 11, 1633-1644.
- Robledo, G. L., Urcelay, C., Domínguez, L., & Rajchenberg, M. (2006) Taxonomy, ecology and biogeography of polypores (*Basidiomycetes*) from Argentinian *Polylepis* woodlands. *Canadian Journal of Botany*, 84, 1561-1572.
- Robledo, G. L. & Renison, D. L. (2010). Wood-decaying polypores in the mountains of central Argentina in relation to *Polylepis* forest structure and altitude. *Fungal Ecology*, 3, 178-184.
- Ryvarden, L. & Johansen, I. (1980). *A preliminary polypore flora of East Africa*. Oslo: Fungiflora.
- Ryvarden, L. (1991). Genera of polypores, nomenclature and taxonomy. *Synopsis Fungorum*, 5, 1-363.
- Ryvarden, L. (2004). Neotropical Polypores. Part 1. *Synopsis Fungorum*, 19, 69-103.
- Schaeffer-Novelli, Y., Cintrón-Molero, G. R., Andaime, R., & Camargo, T. M. (1990). Variability of the Mangrove Ecosystems along the Brazilian coast. *Estuaries*, 13(2), 204-218.
- Schnittler, M. & Stephenson, S. L. (2000). Myxomycete biodiversity in four different forest types in Costa Rica. *Mycologia*, 92(4), 626-637.
- Silva, M. A. B., Bernini, E., & Carmo, T. M. S. (2005). Características estruturais de bosques de mangue do estuário do rio São Mateus, ES, Brasil. *Acta Botanica Brasílica*, 19(3), 465-471.
- Sothão, H. M. P., Bononi, V. L. R., & Figueiredo, T. S. (1991). Basidiomycetes de manguezais da Ilha de Maracá, Amapá, Brasil. *Boletim do Museu Paraense Emílio Goeldi, Série Botânica*, 7(1), 109-114.
- Trierveiler-Pereira, L., Santos P. J. P., & Baseia, I. G. (2013). Ecological aspects of epigeous gasteromycetes (Agaricomycetes, Basidiomycota) in four remnants of the Brazilian Atlantic Forest. *Fungal Ecology*, 6, 471-476.
- Urcelay, C. & Robledo, G. (2004). Community structure of polypores (*Basidiomycota*) in Andean Alder wood in Argentina: Functional groups among wood-decay fungi? *Austral Ecology*, 29, 471-476.
- Webster, J. & Weber, R. W. S. (2007). *Introduction to Fungi*. (3rd ed.). Cambridge: Cambridge University Press.
- Yamashita, S., Hattori, T., Momose, K., Nakagawa, M., Aiba, M., & Nakashizuka, T. (2008). Effects of Forest Use on Aphyllophoraceous Fungal Community Structure in Sarawak, Malaysia. *Biotropica*, 40(3), 354-362.